

Abuse, Violence and/or Intimate Partner Violence – HM 11

DEFINITION

Intimate partner violence (IPV) is abuse or aggression that occurs in a romantic relationship, “Intimate partner” refers to both current, former, and dating partner. CDC identifies 4 types of IPV – physical violence, sexual violence, stalking and psychological aggression. Psychological aggression is estimated to be the most common form of IPV. A pattern of coercive behavior used to gain power and control over another person through fear and intimidation. All individuals are at risk for abuse and/or violence regardless of age, race or socioeconomic status.

SUBJECTIVE

May include:

1. Client may describe episodes of physical, sexual, psychological or verbal abuse directed at themselves or others.
2. Client may present with injuries inconsistent with the history described.
3. Client may have a history of frequent visits to health care providers where they present with multiple injuries or vague somatic complaints (i.e., headaches, GI complaints, fatigue, sleeplessness, sexual dysfunction, chest pain, palpitations, allergic skin reactions, musculoskeletal aches, anxiety).
4. Client may be accompanied by an over-protective partner who does not want the client to be left alone with the health care provider.
5. Client may report history of:
 - a. Missed appointments or presenting for treatment days after an injury.
 - b. Alcohol or substance use in self or partner.
 - c. Eating disorders, depression, panic attacks, suicidal ideation or suicide attempt(s).
 - d. Pre-term labor, low birth weight infant, or miscarriage in previous pregnancies.
6. Strangulation risks may be present in suicide attempts, domestic violence, sports injuries, consensual and non-consensual sexual activity, martial arts and law enforcement activities.

OBJECTIVE

May include:

1. Client may appear restless, angry, defensive, tearful, evasive, or anxious. May also exhibit an inappropriate affect or avoid eye contact.
2. Client may present with:
 - a. Patchy alopecia
 - b. Cigarette burns, human bites, multiple injuries and/or bruises in various stages of healing, wounds to the face, head, neck, breasts, or abdomen, wounds from a knife or firearm. (See Century Code 43-17-41, Duty of Physicians and Others to Report Injury.
 - c. Foreign objects in ear, nose, vagina or rectum
 - d. Conditions associated with stress (i.e., hypertension, obesity, weight loss, GI ulcer)
 - e. Evidence of sexual abuse (i.e., lacerations on breast, labia, urethra, perineum/anal area)
 - f. Gynecological problems (i.e., frequent vaginal and UTI's, pelvic pain, STI's)
 - g. Signs of possible strangulation (i.e., conjunctival or facial petechia, neck abrasions, edema, ligature marks, hoarseness or stridor)

LABORATORY

As indicated by physical findings

ASSESSMENT

Abuse and/or violence

Effective Date: 12/1/2023

Last Reviewed: 10/24/2023

Next Scheduled Review: 10/1/2024

PLAN

May include:

1. Screen clients regardless of age, race, socioeconomic or marital status, as indicated. Discuss reporting requirements prior to screening.
2. Speak to the client alone.
3. Assure the client of confidentiality.
4. Referral to a local crisis program, as indicated. May arrange for crisis counselor to come to Family Planning Clinic. See [Find Help | CAWS North Dakota](#)
5. Provide client referral information to community resources such as legal, law enforcement, shelter, financial and counseling services, as indicated.
6. Use caution if prescribing sedatives, tranquilizers or antidepressants. (They could be used by the client in a suicide attempt.)
7. Document findings in a clear, precise and comprehensive fashion using diagrams, measurements and photographs (if available and if client consents).

CLIENT EDUCATION

1. Inform all clients that abuse, and violence is not a normal part of relationships.
2. Educate client about the cycle of violence and explain that abuse/violent episodes will likely increase in frequency and severity without intervention.

CONSULT / REFER TO PHYSICIAN

1. Medical consultation/referral as appropriate for treatment of injuries.
2. Mental health or substance abuse consultation/referral as appropriate.
3. Urgent referral for report of strangulation assault for evaluation of unknown acute injury and uncertain prognosis.

REFERENCES

1. North Dakota Council on Abused Women's Services and Coalition Against Sexual Assault in North Dakota [Find Help | CAWS North Dakota](#)
2. National Consensus Guidelines-Futures Without Violence. [NCG_final.pdf \(futureswithoutviolence.org\)](#)
3. North Dakota Family Planning Program website. [Home - ND Family Planning \(ndhealth.gov\)](#)
4. National Domestic Violence Hotline [Domestic Violence Support | The National Domestic Violence Hotline \(thehotline.org\)](#)
5. AHRQ. Intimate Partner Violence Screening. Best Screening Practices (tools identified on website). [Intimate Partner Violence Screening | Agency for Healthcare Research and Quality \(ahrq.gov\)](#)
6. <https://www.ncbi.nlm.nih.gov/books/NBK459192/> Strangulation injuries 6/2022

Effective Date: 12/1/2023

Last Reviewed: 10/24/2023

Next Scheduled Review: 10/1/2024